

# FIFRA Section 24(c) Special Local Need

GROUP 14 HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OKLAHOMA

## CHATEAU®

HERBICIDE SW

FOR USE IN ROW MIDDLES OF TRANSPLANTED PEPPERS  
(BELL AND NON-BELL PEPPERS)

EPA Reg. No. 59639-99  
EPA SLN No. OK-11-0003

Active Ingredient	By Wt.
Flumioxazin* .....	51%
Other Ingredients .....	49%
Total .....	100%

\*2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isindole-1,3(2H)-dione

*Chateau* Herbicide SW is a water dispersible granule containing 51% active ingredient.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Persons using this product must comply with all applicable directions, restrictions, Worker Protections Standard requirements and precautions found on the label of the federally registered product upon which this registration is based.

This labeling must be in the possession of the user at the time of pesticide application.

**READ ENTIRE LABEL AND BOOKLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Application Timing	Use Rate Per Application	Special Use Precautions
Pre-Transplant	2 oz/A	Use up to 2 oz/A pre-transplant of <i>Chateau</i> Herbicide SW plus 2 oz/A post-transplant. Not to exceed 4 oz/A per year
Post-Transplant	2 oz/A	

### RESTRICTIONS AND LIMITATIONS

- Apply *Chateau* Herbicide SW only in row middles between rows of pepper (bell and non-bell).
- Spray must remain in row middles between rows of pepper with no contact with the existing crop by applying with either a hooded or shielded sprayer.
- May be applied to row middles pre-transplant or post-transplant
- Do not apply more than 4 oz of *Chateau* Herbicide SW per acre during a single application.
- Injury can occur if soil particles treated with *Chateau* Herbicide SW contact the crop.
- Rainfall or irrigation after application is required prior to transplanting.

### TIMING TO CROP

*Chateau* Herbicide SW may be applied at 4 oz per acre as a shielded or hooded spray to row middles before or after transplanting.

### WEED CONTROL AND TANK MIXING

*Chateau* Herbicide SW provides preemergence residual control of the weeds listed in Table 1 of the container label as well as to assist in the postemergence control of emerged weeds. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For control of emerged weeds, tank-mix *Chateau* Herbicide SW with paraquat, Aim, glyphosate, or other registered burndown herbicide. Refer to tank mix partner label for recommended rates and application parameters.

**Table 1. Broadleaf Weeds Controlled\* by Residual Activity of CHATEAU HERBICIDE SW**

BROADLEAF WEED SPECIES				
SECTION A				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	RATE
Carpetweed	<i>Mollugo verticillata</i>	Up to 5%	All Soil Types	2 oz/A
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>			
Field Pennycress	<i>Thlaspi arvense</i>			
Florida Pusley	<i>Richardia scabra</i>			
Henbit	<i>Lamium amplexicaule</i>			
Lambsquarters, Common	<i>Chenopodium album</i>			

**\*Refer to Container Label for additional weeds controlled.**

**Table 1. (Cont.) Broadleaf Weeds Controlled\* by Residual Activity of CHATEAU HERBICIDE SW**

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	RATE
Little Mallow	<i>Malva parviflora</i>	Up to 5%	All Soil Types	2 oz/A
Marestail/Horseweed	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria maritima</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds				
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane, Common	<i>Portulaca oleracea</i>			
Radish, Wild	<i>Raphanus raphanistrum</i>			
Redmaids	<i>Calandrinia ciliata</i> var. <i>menziesii</i>			
Shepherd's-purse	<i>Capsella bursa-pastoris</i>			
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>			
Sowthistle, Prickly	<i>Sonchus asper</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Venice Mallow	<i>Hibiscus trionum</i>			

**\*Refer to Container Label for additional weeds controlled.**

FIFRA §24(c) Registrant: Valent USA Corporation  
P.O. Box 8025  
Walnut Creek, CA 94596

*Chateau* is a registered trademark of Valent U.S.A. Corporation

**This label is valid through December 31, 2016 or until otherwise amended, withdrawn, cancelled or suspended.**